

HOMOLOGY OF DIHEDRAL QUANDLES

MACIEJ NIEBRZYDOWSKI AND JÓZEF H. PRZYTICKI

ABSTRACT. We solve the conjecture by R. Fenn, C. Rourke and B. Sanderson that the rack homology of dihedral quandles satisfies $H_3^R(R_p) = \mathbb{Z} \oplus \mathbb{Z}_p$ for p odd prime ([Oht], Conjecture 5.12). We also show that $H_n^R(R_p)$ contains $\mathbb{Z}_p$ for $n \geq 3$. Furthermore, we show that the torsion of $H_n^R(R_3)$ is annihilated by 3. We also prove that the quandle homology $H_4^Q(R_p)$ contains $\mathbb{Z}_p$ for p odd prime. We conjecture that for $n > 1$ quandle homology satisfies: $H_n^Q(R_p) = \mathbb{Z}_p^{f_n}$, where f_n are “delayed” Fibonacci numbers, that is, $f_n = f_{n-1} + f_{n-3}$ and $f(1) = f(2) = 0, f(3) = 1$. Our paper is the first step in approaching this conjecture.

CONTENTS

1. Definitions and Preliminary facts	1
2. Homological operations	4
3. The proof of the Fenn-Rourke-Sanderson conjecture	9
4. 3 annihilates the torsion of $H_n^R(R_3)$	10
5. $H_4^Q(R_p)$ is nontrivial for p odd prime	14
6. Future directions	17
7. Appendix 1	18
8. Appendix 2	19
References	20

1. DEFINITIONS AND PRELIMINARY FACTS

In 1942, Mituhisa Takasaki introduced the notion of kei [Tak] as an abstraction of the notion of symmetric transformation. He analysed, in particular, finite keis corresponding to symmetries of regular polygons (today they are called dihedral quandles). Kei is an abstract algebra $(Q, *)$ with a binary operation $* : Q \times Q \rightarrow Q$ satisfying the conditions:

- (i) $a * a = a$, for any $a \in Q$;
- (ii) $(a * b) * b = a$, for any $a, b \in Q$;

Date: November 25, 2006.

1991 *Mathematics Subject Classification.* Primary 55N35; Secondary 18G60, 57M25.

Key words and phrases. dihedral quandle, Alexander quandle, rack, quandle homology, homological operations.

- (iii) $(a * b) * c = (a * c) * (b * c)$, for any $a, b, c \in Q$ (the right distributivity property).

If the second condition is relaxed to the following condition:

(ii') for every $b \in Q$ the map $*_b : Q \rightarrow Q$, given by $*_b(a) = a * b$, is a bijection, then $(Q, *)$ is called a quandle (the name coined by Joyce [Joy]). A kei is an involutive $((*_b)^2 = Id)$ quandle. If the first condition is omitted, the abstract algebra we obtain is called a rack (J. H. Conway in his 1959 correspondence with G. C. Wraith suggested *wrack* which was changed to rack¹ in [F-R]).

Takasaki was considering keis associated to abelian groups, that is, $kei(G)$ satisfies $a * b = 2b - a$, for $a, b \in G$. The k -dihedral quandle, denoted R_k , is isomorphic to $kei(\mathbb{Z}_k)$. Generalization of a kei of abelian group to Alexander quandle of $Z[t, t^{-1}]$ -module can be traced back to R. Fox and his suggestion that arcs of the link diagram can be colored using polynomials. The Alexander quandle $Alex(M)$ can be associated to every $Z[t, t^{-1}]$ module M , by taking the quandle operation to be $a * b = (1 - t)a + tb$ for any $a, b \in M$.

Rack homology and homotopy theory were first defined and studied in [FRS], and a modification to quandle homology theory was given in [CJKLS] to define knot invariants in a state-sum form (so-called cocycle knot invariants).

We recall here the definition of rack, degenerate and quandle homology after [CKS].

Definition 1.1. (i) For a given rack X , let $C_n^R(X)$ be the free abelian group generated by n -tuples $(x_1, x_2, \dots, x_n)$ of elements of X ; in other words, $C_n^R(X) = \mathbb{Z}X^n = (\mathbb{Z}X)^{\otimes n}$. Define a boundary homomorphism $\partial : C_n^R(X) \rightarrow C_{n-1}^R(X)$ by:

$$\partial(x_1, x_2, \dots, x_n) = \sum_{i=2}^n (-1)^i ((x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) - (x_1 * x_i, x_2 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)).$$

$(C_*^R(X), \partial)$ is called a rack chain complex of X .

- (ii) Assume that X is a quandle. Then we have a subchain complex $C_n^D(X) \subset C_n^R(X)$, generated by n -tuples $(x_1, \dots, x_n)$ with $x_{i+1} = x_i$ for some i . The subchain complex $(C_n^D(X), \partial)$ is called a degenerated chain complex of a quandle X .

¹From [F-R]: The earliest work on racks (known to us) is due to Conway and Wraith [correspondence, 1959]... They used the name *wrack* for the concept and we have adopted this name, not merely because it is the oldest name, but also because it is a simple English word which (to our knowledge) has no other mathematical meaning. We have however chosen the more common spelling. Rack is used in the same sense as in the phrase “rack and ruin”. The context of Conway and Wraith’s work is the conjugacy operation in a group and they regarded a rack as the wreckage of a group left behind after the group operation is discarded and only the notion of conjugacy remains.

- (iii) The quotient chain complex $C_n^Q(X) = C_n^R(X)/C_n^D(X)$ is called the quandle chain complex. We have the short exact sequence of chain complexes:

$$0 \rightarrow C_n^D(X) \rightarrow C_n^R(X) \rightarrow C_n^Q(X) \rightarrow 0.$$

- (iv) The homology of rack, degenerate, and quandle chain complexes are called rack, degenerate, and quandle homology, respectively. We have the long exact sequence of homology of quandles:

$$\dots \rightarrow H_n^D(X) \rightarrow H_n^R(X) \rightarrow H_n^Q(X) \rightarrow H_{n-1}^D(X) \rightarrow \dots$$

- (v) For an abelian group G , define the chain complex

$C_*^Q(X; G) = C_*^Q \otimes G$, with $\partial = \partial \otimes id$. The groups of cycles and boundaries are denoted respectively by $\ker(\partial) = Z_n^Q(X; G) \subset C_n^Q(X; G)$ and $Im(\partial) = B_n^Q(X; G) \subset C_n^Q(X; G)$. The n th quandle homology group of a quandle X with coefficient group G is defined as

$$H_n^Q(X; G) = H_n(C_*^Q(X; G)) = Z_n^Q(X; G)/B_n^Q(X; G).$$

R. Litherland and S. Nelson [L-N] proved that the short exact sequence from (iii) splits respecting the boundary maps.

The splitting map, $\alpha : C_n^Q(X) \rightarrow C_n^R(X)$, is given by:

$$\alpha(x_1, x_2, x_3, \dots, x_n) = (x_1, x_2 - x_1, x_3 - x_2, \dots, x_n - x_{n-1}),$$

where, in our notation, $(x_1, x_2 - x_1, x_3 - x_2, \dots, x_n - x_{n-1}) = x_1 \otimes (x_2 - x_1) \otimes (x_3 - x_2) \otimes \dots \otimes (x_n - x_{n-1}) \in C_n^R(X)$. In particular, α is a chain complex monomorphism and $H_n^R(X) = H_n^D(X) \oplus \alpha_*(H_n^Q(X))$.

Free part of homology of quandles ($free(H_*(X))$) was computed in [L-N, E-G] (lower bounds for Betti numbers were given in [CJKS]). In particular, it was shown there that for dihedral quandle R_k , k odd, we have:

Theorem 1.2. (i) [L-N, E-G]

$$free(H_n^R(R_k)) = \mathbb{Z}$$

(ii) [L-N, E-G]

$$free(H_n^Q(R_k)) = \begin{cases} \mathbb{Z} & \text{for } n = 1 \\ 0 & \text{otherwise} \end{cases}$$

Useful information concerning torsion of homology of racks and quandles was obtained in [L-N, Moc]. In particular, it was shown that:

Theorem 1.3. (i) [L-N, Moc] Torsion of $H_n^R(R_k)$ is annihilated by k^n .

(ii) [Moc] The third cohomology $H_Q^3(R_p; \mathbb{Z}_p) = \mathbb{Z}_p$ for p odd prime.

(iii) [L-N] For a quandle X , let $\mathcal{O}_X$ denote the set of orbits of X with respect to the action of X on itself by the right multiplication (e.g. $|\mathcal{O}_{R_k}|$ is 1 for odd k and 2 for an even k). Then

$$H_2^R(X) \cong H_2^Q(X) \oplus \mathbb{Z}\mathcal{O}_X,$$

$$H_3^R(X) \cong H_3^Q(X) \oplus H_2^Q(X) \oplus \mathbb{Z}\mathcal{O}_X^2.$$

In particular, $H_3^R(R_k) \cong H_3^Q(R_k) \oplus \mathbb{Z}$ for k odd.

We devote this paper to computation of rack and quandle homology of odd dihedral quandles. One of the results of this paper is the solution of the conjecture by Fenn, Rourke and Sanderson, listed as Conjecture 5.12 in Ohtsuki's problem list [Oht].

Conjecture 1.4 (R. Fenn, C. Rourke, B. Sanderson).

$H_3^R(R_p) \cong \mathbb{Z} \oplus \mathbb{Z}_p$ for p odd prime.

We prove Conjecture 1.4 in Section 3. We also prove the generalization of Theorem 1.3(i) for $k = 3$, namely, we show that 3 annihilates the torsion of $H_n^R(R_3)$ for any n . We also propose the general conjecture on the structure of quandle homology $H_n^Q(R_k)$.

Conjecture 1.5. For a prime number p , $\text{tor} H_n^Q(R_p) = \mathbb{Z}_p^{f_n}$, where $\{f_n\}$ are “delayed” Fibonacci numbers, that is,

$$f_n = f_{n-1} + f_{n-3}, \text{ and } f(1) = f(2) = 0, f(3) = 1.$$

We verified the conjecture by GAP [GAP4] calculation for $p = 3$ and $n \leq 12$, $p = 5$ and $n \leq 6$, and for $p = 7$ and $n \leq 4$.

We also computed² that $H_3^Q(R_9) = H_4^Q(R_9) = \mathbb{Z}_9$, so there is a possibility that Conjecture 1.5 may hold for some non-prime odd k (e.g., powers of odd prime numbers). One cannot also exclude that it holds for any odd k .

2. HOMOLOGICAL OPERATIONS

In this section we construct two chain maps raising index of chain groups of a rack (and quandle) by one and two, respectively. We use these maps in Sections 3 and 4 to prove our main results on homology of racks and quandles. We expect to find one more homology operation, from $H_n(R_k)$ to $H_{n+4}(R_k)$, but it is still an open problem. In this paper, we are mostly interested in homology of dihedral quandles, but sometimes we formulate more general results, if it doesn't make proofs much longer.

In general, our homological operations involve the group homomorphism $h_u : C_n^R(X) \rightarrow C_{n+j}^R(X)$, for $u \in \mathbb{Z}X^j$, given by $h_u(w) = (w, u)$. This map usually is not a chain map, unless we choose a special u and/or consider quandles satisfying some special conditions (e.g., as in Lemma 2.5).

The first homological operation is related to the group homomorphism $h_a : C_n^R(X) \rightarrow C_{n+1}^R(X)$, given by $h_a(w) = (w, a)$, for any $a \in X$, and $w \in X^n$. This map is not a chain map, so we need to symmetrize it with respect to another map $*_a : C_n^R(X) \rightarrow C_n^R(X)$ given by $*_a(w) = w * a$, for any $w \in X^n$, or more precisely, $*_a(x_1, \dots, x_n) = (x_1, \dots, x_n) * a =$

² The cases $p = 3$, $n = 12$, $p = 5$, $n = 6$ and $p = 9$, $n = 4$ were computed with the help of A. Shumakovitch and Norbert A'Campo, who provided computer power of his supercomputer located at Basel University, Switzerland.

$(x_1 * a, \dots, x_n * a)$ (the map $*_a$ for $n = 1$ is exactly the map used in Condition (ii') of a quandle). In other words, we consider a function $h'_a = h_a + *_a h_a$. The basic properties of these maps are described in the following proposition:

Proposition 2.1.

- (i) For any rack X , and $a \in X$, the map $*_a$ is a chain map chain homotopic to the identity.
- (ii) If X is a kei (involutive quandle), then³ $h'_a = h_a + *_a h_a$ is a chain map.
- (iii) If X is a quandle, then $h_a(C_n^D(X)) \subset C_{n+1}^D(X)$, $*_a(C_n^D(X)) \subset C_n^D(X)$. Therefore, the maps $h_a, h'_a : C_n^Q(X) \rightarrow C_{n+1}^Q(X)$ are well defined.
- (iv) If a and b are in the same orbit of X , then h'_a and h'_b induce the same map on homology, that is, $(h'_a)_* = (h'_b)_* : H_n^R(X) \rightarrow H_{n+1}^R(X)$.

Proof. (i) $*_a$ is a chain map, because⁴

$$\begin{aligned}
 d(*_a(x_1, \dots, x_n)) &= d(x_1 * a, \dots, x_n * a) = \sum_{i=2}^k (-1)^i ((x_1 * a, \dots, x_{i-1} * a, x_{i+1} * a, \dots, x_n * a) \\
 &\quad - (x_1 * a * (x_i * a), \dots, x_{i-1} * a * (x_i * a), x_{i+1} * a, \dots, x_n * a)) = \\
 &\sum_{i=2}^k (-1)^i ((x_1 * a, \dots, x_{i-1} * a, x_{i+1} * a, \dots, x_n * a) - (x_1 * x_i * a, \dots, x_{i-1} * x_i * a, x_{i+1} * a, \dots, x_n * a)) \\
 &= *_a \left(\sum_{i=2}^k (-1)^i ((x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) - (x_1 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)) \right) = \\
 &\quad *_a(d(x_1, \dots, x_n)).
 \end{aligned}$$

The homomorphism $(-1)^{n+1} h_a : C_n^R(X) \rightarrow C_{n+1}^R(X)$ is a chain homotopy between Id and $*_a$ chain maps. Namely:

$$\begin{aligned}
 d((-1)^{n+1} h_a(x_1, \dots, x_n)) &= \\
 (-1)^{n+1} ((d(x_1, \dots, x_n), a) + (-1)^{n+1} ((x_1, \dots, x_n) - (x_1 * a, \dots, x_n * a))) &= \\
 -(-1)^n h_a(d(x_1, \dots, x_n)) + (Id - *_a)(x_1, \dots, x_n),
 \end{aligned}$$

as needed.

$$\begin{aligned}
 \text{(ii) } dh'_a &= d(h_a + *_a h_a) = dh_a + (dh_a) * a = h_a d + (-1)^{n+1} (Id - *_a) + (h_a d + \\
 &\quad (-1)^{n+1} (Id - *_a)) * a = h_a d + (h_a * a) d = h'_a d.
 \end{aligned}$$

(iii) It follows from the definition of rack, degenerate and quandle chain complex of X .

³To be more general, we have to assume that a satisfies the “n-condition”, that is, $x * a * a * \dots * a = x$ for any x , and take $h'_a = h_a + *_a h_a + *_a *_a h_a + \dots + *_a(\dots(*_a h_a))$. For example, for quandle S_4 , $h'_a = h_a + *_a h_a + *_a *_a h_a$. Then h'_a is a chain map. Notice that if X is finite, then any $a \in X$ satisfies n condition for some n .

⁴ We use a standard convention for products in non-associative algebras, called the left normed convention, that is, whenever parentheses are omitted in a product of elements $a_1, a_2, \dots, a_n$ of Q then $a_1 * a_2 * \dots * a_n = ((\dots((a_1 * a_2) * a_3) * \dots) * a_{n-1}) * a_n$ (left association), for example, $a * b * c = (a * b) * c$.

(iv) It suffices to consider the case, when there is $x \in X$, such that $b = a * x$. Notice, that

$$\begin{aligned} *_x(h'_a(w)) &= (h'_a(w)) * x = ((w + w * a), a) * x = ((w * x + (w * a) * x), b) = \\ &= ((w * x + (w * x) * b), b) = h'_b(w * x) = h'_b(*_x(w)). \end{aligned}$$

On the other hand, if w is a cycle then, by (i), $*_x(w)$ is a homologous cycle. Therefore, $h'_b(*_x(w))$ and $h'_b(w)$ are homologous by (ii). Similarly, $*_x(h'_a(w))$ and $h'_a(w)$ are homologous. Therefore, $h'_b(w)$ and $h'_a(w)$ are homologous. $\square$

Next, we prove that under certain assumptions that hold for odd dihedral quandles, the map $(h'_a)_*$ is a monomorphism on rack homology.

Definition 2.2. We say that a rack X satisfies:

- the property (1), if for any elements $x, y, a \in X$, $*_x(a) = *_y(a)$ implies that the maps $*_x$ and $*_y$ are equal (i.e., $z * x = z * y$, for any $z \in X$);
- the property (2), if for any elements $x, y, x', y', a \in X$, from $a * x * x' = a * y * y'$ follows that the maps $*_{x'} * x$ and $*_{y'} * y$ are equal (i.e., $z * x' * x = z * y' * y$, for any $z \in X$);
- the quasigroup property, if for any $a, b \in X$, the equation $a * x = b$ has exactly one solution.

An example of racks satisfying properties (1) and (2) is algebraically connected (i.e., with one orbit) Alexander quandles. Indeed, in such case we can write $*_x(a) = *_y(a)$ as $(1-t)x + ta = (1-t)y + ta$, so $(1-t)x = (1-t)y$, and that implies property (1).

Similarly, $a * x * x' = a * y * y'$ is equivalent to

$$(1-t)(x' + tx) + t^2a = (1-t)(y' + ty) + t^2a,$$

and further to

$$(1-t)(x' + tx) = (1-t)(y' + ty),$$

from which property (2) follows.

The quasigroup property is stronger than property (1) and an Alexander quandle M possesses this property only if $(1-t)$ does not annihilate non-zero elements in the module M , and if division by $1-t$ is possible. Odd dihedral quandles satisfy all above properties.

Theorem 2.3. Let a kei X be a quasigroup satisfying property (2) of Definition 2.2. Then $(h'_a)_* : H_n^R(X) \rightarrow H_{n+1}^R(X)$ has the following property.

There is a map $\bar{h}_* : H_{n+1}^R(X) \rightarrow H_n^R(X)$ such that $\bar{h}_*(h'_a)_* = 4Id$. In particular, if $H_n^R(X)$ has no element of order 2, then $(h'_a)_*$ is a monomorphism.

Proof. From the Proposition 2.1(ii), we know that $h'_a = h_a + *_a h_a$ is a chain map. Let $\bar{h}_a : C_{n+1}^R(X) \rightarrow C_n^R(X)$ be given by

$$\bar{h}_a(x_1, \dots, x_n, x_{n+1}) = (x'_1, \dots, x'_n),$$

where $(x'_1, \dots, x'_n)$ is uniquely determined by the equality (with unique x)

$$(x_1, \dots, x_n, x_{n+1}) * x = (x'_1, \dots, x'_n, a).$$

We will show that the map $\bar{h}'_a = \bar{h}_a + *_a \bar{h}_a$ is a chain map. We have

$$\begin{aligned} d(\bar{h}'_a(x_1, \dots, x_n, x_{n+1})) &= d((x_1, \dots, x_n) * x + (x_1, \dots, x_n) * x * a) = \\ &= (d(x_1, \dots, x_n)) * x + (d(x_1, \dots, x_n)) * x * a. \end{aligned}$$

We also have

$$\begin{aligned} \bar{h}'_a(d(x_1, \dots, x_n, x_{n+1})) &= \bar{h}'_a(d(x_1, \dots, x_n), x_{n+1}) + \\ &= (-1)^{n+1}(\bar{h}_a(x_1, \dots, x_n) - \bar{h}_a((x_1, \dots, x_n) * x_{n+1}) + \\ &= (\bar{h}_a(x_1, \dots, x_n)) * a - (\bar{h}_a((x_1, \dots, x_n) * x_{n+1})) * a). \end{aligned}$$

To show that $\bar{h}'_a$ is a chain map, we need equalities:

$$\bar{h}_a(x_1, \dots, x_n) = (\bar{h}_a((x_1, \dots, x_n) * x_{n+1})) * a,$$

and

$$(\bar{h}_a(x_1, \dots, x_n)) * a = \bar{h}_a((x_1, \dots, x_n) * x_{n+1}).$$

Notice, that these equalities are equivalent when X is a kei.

In order to prove them, we need all assumptions stated in the theorem.

Let x and y be such that $x_n * x = a$ and $x_n * x_{n+1} * y = a$. Therefore, $x_n * x * a = x_n * x_{n+1} * y$. It follows, by property (2), that the maps $*_a *_x$ and $*_y *_x$ are equal on the entire X , and the required equalities follow.

Once we know that $\bar{h}'_a$ is a chain map, we can finish the proof by noticing that

$$\bar{h}'_a h'_a(w) = w + w * a + w * a + w * a * a = 2(w + w * a).$$

Therefore, on homology, $(\bar{h}'_a h'_a)_* = 4Id$.

From the Theorem 1.3(i) follows that, for k odd, $H_n^R(R_k)$ can only have odd torsion. Therefore, for k odd, $(h'_a)_* : H_n^R(R_k) \rightarrow H_{n+1}^R(R_k)$ is a monomorphism. $\square$

Corollary 2.4. *Let X be a kei satisfying the conditions of Theorem 2.3.*

- (i) *If $\text{tor} H_n^R(X)$ is annihilated by N , then $\text{tor} H_{n+1}^R(X)$ is annihilated by $4|X|N$.*
- (ii) *If $\text{tor} H_n^R(R_k)$, k odd, is annihilated by N then $\text{tor} H_{n+1}^R(R_k)$ is annihilated by kN .*
- (iii) *For k odd, $\text{tor} H_3^R(R_k)$ is annihilated by k .*

Proof. (i) Consider the (symmetrizer) chain map $s_{n+1} : C_{n+1}(X) \rightarrow C_{n+1}(X)$ given by

$$s_{n+1}(u) = \sum_{x \in X} (u * x).$$

By Proposition 2.1(i) the map $(s_{n+1})_* : H_{n+1}^R(X) \rightarrow H_{n+1}^R(X)$ is equal to $|X|Id$. Next, consider the composition

$$s_{n+1} h'_a \bar{h}'_a : C_{n+1}(X) \rightarrow C_{n+1}(X).$$

To find $s_{n+1}h'_a\bar{h}'_a(w, x_{n+1})$, we assume that $x_{n+1} * x = a$ (so that we also know that $x_{n+1} = a * x$). Then:

$$\begin{aligned} s_{n+1}h'_a\bar{h}'_a(w, x_{n+1}) &= s_{n+1}h'_a(w * x + w * x * a) = s_{n+1}(2((w * x, a) + (w * x * a, a))) = \\ 2 \sum_{y \in X} ((w * x * y, a * y) + (w * x * a * y, a * y)) &= 2 \sum_{y \in X} ((w, x_{n+1}) * y * a + (w, x_{n+1}) * y). \end{aligned}$$

In order to prove the last equality, we need to show that:

- (1) $\sum_{y \in X} (w * x * y, a * y) = \sum_{y \in X} (w, x_{n+1}) * y * a$;
- (2) $\sum_{y \in X} (w * x * a * y, a * y) = \sum_{y \in X} (w, x_{n+1}) * y$.

The left hand side of (1) is equal to

$$\sum_{y \in X} (w * x * y, a * y) = \sum_{y \in X} (w * x, a) * y = \sum_{y \in X} (w, x_{n+1}) * x * y.$$

Now, the set $\{x_{n+1} * x * y : y \in X\}$ is the set of all elements of X , as is the set $\{x_{n+1} * y * a : y \in X\}$. Furthermore, by Property (2) of Definition 2.2, if $x_{n+1} * x * y = x_{n+1} * y' * a$ for some y and y' , then $w * x * y = w * y' * a$ for any $w \in C_n(X)$. Thus, the equality (1) holds.

To prove (2), we notice that:

$$\begin{aligned} \sum_{y \in X} (w * x * a * y, a * y) &= \sum_{y \in X} (w * x, a) * (y * a) * a = \\ \sum_{y \in X} (w, x_{n+1}) * x * (y * a) * a &= \sum_{y \in X} (w, x_{n+1}) * x * y * a. \end{aligned}$$

By a similar argument as in (1), for every $y \in X$ there exists $y' \in X$ such that $x_{n+1} * x * y * a = x_{n+1} * y' * a$, i.e., $x_{n+1} * x * y = x_{n+1} * y' * a$. It follows that $w * x * y = w * y' * a$ (and $w * x * y * a = w * y' * a$) for any $w \in C_n(X)$, and that finishes the proof of equality (2).

By Proposition 2.1(i) the map $(s_{n+1}h'_a\bar{h}'_a)_* = 4|X|Id$. Therefore, if $torH_n^R(X)$ is annihilated by N , then $torH_{n+1}^R(X)$ is annihilated by $4|X|N$.

(ii) By Theorem 1.3(i), $H_n^R(R_k)$, k odd, has no even torsion. Thus, if $torH_n^R(X)$ is annihilated by N , then $torH_{n+1}^R(X)$ is annihilated by kN .

(iii) As given in [CJKS] (Example 3.10), $torH_2^R(R_k) = 0$ for k odd. Thus, $torH_3^R(R_k)$ is annihilated by k . $\square$

The second homological operation, of degree two, is defined on somehow restricted class of racks (and quandles), but including dihedral quandles and Burnside keis (see footnote 5). Let X be any rack with elements $x_0, x_1, \dots, x_{k-1}$, satisfying the “cyclic” Fibonacci relation⁵ modulo k . That is, $x_2 = x_0 * x_1$, $x_3 = x_1 * x_2, \dots, x_0 = x_{k-2} * x_{k-1}$ and $x_1 = x_{k-1} * x_0$.

⁵If X is a kei, then by eliminating all generators except x_0 and x_1 , we obtain relations $x_0 = (((\dots x_0) * x_1) * x_0) * x_1$ and $x_1 = (((\dots x_1) * x_0) * x_1) * x_0$, with k letters on the right hand side of each equation. They were studied in [N-P] and led to *Burnside Keis* which are important in analysis of rational moves on links and tangles.

Let $s = s(x_0, x_1) = \sum_{i=0}^{k-1} (x_i, x_{i+1})$, where indices are taken modulo k . Then we can use s to define a chain map as follows.

Lemma 2.5.

- (i) The homomorphism $h_s : C_n^R(X) \rightarrow C_{n+2}^R(X)$ given by $h_s(w) = (w, s)$, is a chain map. Recall that $C_{n+2}^R(X) = \mathbb{Z}X^{n+2} = \mathbb{Z}X^n \otimes \mathbb{Z}X^2$ so (w, s) is another notation for $w \otimes s$.
- (ii) If $X = R_k$, then we can take $x_i = i$, that is,

$$s = s(0, 1) = \sum_{i=0}^{k-1} (i, i+1).$$

For an odd k , we can choose any $0 < j < k$, and take $x_0 = 0$, $x_1 = j$, and generally, $x_i = ij$. Let $u_j \in C_2^R(R_k)$ be defined by $u_j = s(0, j)$. Then $h_{u_j} : C_n^R(R_k) \rightarrow C_{n+2}^R(R_k)$ given by $h_{u_j}(w) = (w, u_j)$, is a chain map.

Proof. (i) We notice that

$$\begin{aligned} d(h_s(w)) &= d(w, s) = \\ (dw, s) + \sum_{i=0}^k (-1)^{n+1} ((w, x_{i+1}) - (w, x_i)) - \sum_{i=0}^k (-1)^{n+1} ((w * x_i, x_{i+1}) - (w * x_{i+1}, x_i * x_{i+1})) = \\ h_s(d(w)) - \sum_{i=0}^k (-1)^{n+1} ((w * x_i, x_{i+1}) - (w * x_{i+1}, x_{i+2})) &= h_s(d(w)), \end{aligned}$$

as required.

- (ii) We check that $ij * (ij + j) = 2ij + 2j - ij = ij + 2j$, so the ‘‘cyclic’’ Fibonacci relation is satisfied and we can use part (i). $\square$

We prove, in the next section, that for k odd prime, $h_s : H_1^Q(R_k) \rightarrow H_3^Q(R_k)$ is an epimorphism $\mathbb{Z} \rightarrow \mathbb{Z}_k$. We conjecture the following.

Conjecture 2.6. For an odd prime number k , and $n > 1$, the homomorphism $h_s : H_n^Q(R_k) \rightarrow H_{n+2}^Q(R_k)$ is a monomorphism.

Example 2.7. We checked (using GAP) that $(h_s)_* : H_3^Q(R_3) \rightarrow H_5^Q(R_3)$ is an isomorphism, sending $(0, 1, 0) + (0, 2, 1)$ to $(0, 1, 0, 1, 2) + (0, 1, 0, 2, 0) + (0, 2, 1, 0, 1) + (0, 2, 1, 2, 0)$. We also checked that $(h_s)_* : H_n(R_3) \rightarrow H_{n+2}(R_3)$ is a monomorphism for $n = 4, 5, 6, 7$. Finally, we checked that $(h_s)_* : H_3(R_5) \rightarrow H_5(R_5)$ is an isomorphism.

3. THE PROOF OF THE FENN-ROURKE-SANDERSON CONJECTURE

Recall that $u_j \in C_2^R(R_k)$ is given by $u_j = \sum_{i=0}^{k-1} (i, i+j)$ and that $(0, u_j) = \sum_{i=0}^{k-1} (0, i, i+j)$.

The generator of the third quandle cohomology $H_Q^3(R_p; \mathbb{Z}_p)$, given by T. Mochizuki in [Moc], is of the form

$$\theta(x_1, x_2, x_3) = \frac{(x_2 - x_1)(2x_3^p - x_2^p - (2x_3 - x_2)^p)}{p}.$$

Proposition 3.1. $\theta((0, u_j)) = j^2 \theta((0, u_1)) = -j^2$ in $\mathbb{Z}_p$. In particular, $(0, u_j)$ is a nontrivial cycle in $H_3^Q(R_p)$.

Proof. In the straightforward evaluation of the Mochizuki's 3-cocycle on $(0, u_j)$, we use the easy to check fact that $(p + a)^p \equiv a^p \pmod{p^2}$, for any $a \in \mathbb{Z}_p$, and that $\theta((0, u_j)) = \sum_{i=0}^{p-1} - (i) \frac{(i+2j)^p + i^p - 2(i+j)^p}{p}$. $\square$

To show that $H_3^Q(R_p) = \mathbb{Z}_p$, we use the universal coefficient theorem for cohomology (see [Ha], Theorem 3.2).

Theorem 3.2. *If a chain complex C of free abelian groups has homology groups $H_n(C)$, then the cohomology groups $H^n(C; G)$ of the cochain complex $\text{Hom}(C_n, G)$ are determined by split exact sequences*

$$0 \rightarrow \text{Ext}(H_{n-1}(C), G) \rightarrow H^n(C; G) \rightarrow \text{Hom}(H_n(C), G) \rightarrow 0.$$

We also recall that $\text{Ext}(H, G) = 0$ if H is free.

Now we can complete the proof of the Fenn-Rourke-Sanderson conjecture. First, recall that $H_3^R(R_p) = H_3^Q(R_p) \oplus \mathbb{Z}$, as summarized in Theorem 1.3(iii). Thus, we should show that $H_3^Q(R_p) = \mathbb{Z}_p$. By Mochizuki's result (Theorem 1.3(ii)), $H_3^Q(R_p; \mathbb{Z}_p) = \mathbb{Z}_p$. Thus, from the universal coefficient theorem for cohomology we have:

$$\mathbb{Z}_p = H_3^Q(R_p; \mathbb{Z}_p) = \text{Hom}(H_3^Q(R_p), \mathbb{Z}_p) \oplus \text{Ext}(H_2^Q(R_p), \mathbb{Z}_p) = \text{Hom}(H_3^Q(R_p), \mathbb{Z}_p),$$

as $H_2^Q(R_p) = 0$. Therefore, $H_3^Q(R_p)$ is equal to $\mathbb{Z}$ or $\mathbb{Z}_p^i$, for $i > 0$. But by the Corollary 2.4 and Theorem 1.2(ii), $H_3^Q(R_p)$ is annihilated by p . Therefore, $H_3^Q(R_p) = \mathbb{Z}_p$, what ends the proof.

It follows from the definition of rack and quandle homology, that $H_1^R(R_p)$ and $H_1^Q(R_p)$ are generated by $(0) \in C_1(R_p)$. It follows from above proofs that $(0, u_1) = \sum_{i=0}^{p-1} (0, i, i+1)$ is a generator of $H_3^Q(R_p)$, for p odd prime. Therefore, we conclude that the map $h_s : H_1^Q(R_p) \rightarrow H_3^Q(R_p)$, that we discussed in the previous section, is an epimorphism $\mathbb{Z} \rightarrow \mathbb{Z}_p$.

4. 3 ANNIHILATES THE TORSION OF $H_n^R(R_3)$

In this section we show that 3 annihilates $\text{tor} H_n^R(R_3)$, for $n > 1$. In the proof we use the fact that R_3 is commutative, i.e.,

$$a * b = 2b - a = -b - a = 2a - b = b * a,$$

for any $a, b \in R_3$. However, there is a possibility that our proof can be generalized (provided that the Conjecture 4.3 is true), therefore, we write the proof below in the more general form and we conjecture the following:

Conjecture 4.1. *If q is odd prime, and $n > 1$, then the torsion subgroup of $H_n^R(R_q)$ is annihilated by q .*

Theorem 4.2. *If $q = 3$, and $n > 1$, then $\text{tor}H_n^R(R_q)$ is annihilated by q .*

Proof. Let $x = (x_1, \dots, x_n) \in (R_q)^n$. We define the chain map $f_n^j: C_n^R(R_q) \rightarrow C_n^R(R_q)$ by

$$f_n^j(x) = \begin{cases} x & \text{for } j = 0 \\ \sum_{y \in R_q} (y, \dots, y, x_{j+1}, x_{j+2}, \dots, x_n) & \text{for } 1 \leq j \leq n \\ f_n^n(x) & \text{for } j > n \end{cases}$$

It was shown in [L-N] that f_n^1 is chain homotopic to q times the identity. We prove that f_n^j is chain homotopic to f_n^{j-1} , for $j \geq 2$, and the chain homotopy is given by the formula

$$D_n^j(x) = \begin{cases} \sum_{y \in R_q} (y, \dots, y, x_j, y, x_{j+1}, \dots, x_n) & \text{for } 1 \leq j \leq n \\ 0 & \text{for } j > n \end{cases}$$

We will use standard notation:

$$\partial_n^0(x) = \sum_{i=1}^n (-1)^i (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n);$$

$$\partial_n^1(x) = \sum_{i=1}^n (-1)^{i+1} (x_1 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n).$$

We need to confirm the equality

$$\partial_{n+1} D_n^j(x) + D_{n-1}^j \partial_n(x) = \pm (f_n^j(x) - f_n^{j-1}(x)).$$

Let us write a detailed list of summands appearing on the left hand side of this equation:

$$\partial_{n+1}^0 D_n^j(x) =$$

$$(1) \quad \sum_{i=1}^{j-1} \sum_{y \in R_q} (-1)^i (y, \dots, y, x_j, y, x_{j+1}, \dots, x_n) +$$

$$(2) \quad (-1)^j \sum_{y \in R_q} (y, \dots, y, x_{j+1}, \dots, x_n) +$$

$$(3) \quad (-1)^{j+1} \sum_{y \in R_q} (y, \dots, y, x_j, x_{j+1}, \dots, x_n) +$$

$$(4) \quad (-1)^{j+2} \sum_{y \in R_q} (y, \dots, y, x_j, y, x_{j+2}, \dots, x_n) +$$

$$(5) \quad \sum_{i=j+2}^n \sum_{y \in R_q} (-1)^{i+1} (y, \dots, y, x_j, y, x_{j+1}, \dots, \widehat{x_i}, \dots, x_n)$$

$$\partial_{n+1}^1 D_n^j(x) =$$

$$(6) \quad \sum_{i=1}^{j-1} \sum_{y \in R_q} (-1)^{i+1} (y, \dots, y, x_j, y, x_{j+1}, \dots, x_n) +$$

$$(7) \quad (-1)^{j+1} \sum_{y \in R_q} (y * x_j, \dots, y * x_j, y, x_{j+1}, \dots, x_n) +$$

$$(8) \quad (-1)^{j+2} \sum_{y \in R_q} (y, \dots, y, x_j * y, x_{j+1}, \dots, x_n) +$$

$$(9) \quad (-1)^{j+3} \sum_{y \in R_q} (y * x_{j+1}, \dots, y * x_{j+1}, x_j * x_{j+1}, y * x_{j+1}, x_{j+2}, \dots, x_n) +$$

$$(10) \quad \sum_{i=j+2}^n \sum_{y \in R_q} (-1)^{i+2} (y * x_i, \dots, y * x_i, x_j * x_i, y * x_i, x_{j+1} * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)$$

$$D_{n-1}^j \partial_n^0 =$$

$$(11) \quad \sum_{i=1}^j \sum_{y \in R_q} (-1)^i (y, \dots, y, x_{j+1}, y, x_{j+2}, \dots, x_n) +$$

$$(12) \quad (-1)^{j+1} \sum_{y \in R_q} (y, \dots, y, x_j, y, x_{j+2}, \dots, x_n) +$$

$$(13) \quad \sum_{i=j+2}^n \sum_{y \in R_q} (-1)^i (y, \dots, y, x_j, y, x_{j+1}, \dots, \widehat{x_i}, \dots, x_n)$$

$$D_{n-1}^j \partial_n^1 =$$

$$(14) \quad \sum_{i=1}^j \sum_{y \in R_q} (-1)^{i+1} (y, \dots, y, x_{j+1}, y, x_{j+2}, \dots, x_n) +$$

$$(15) \quad (-1)^{j+2} \sum_{y \in R_q} (y, \dots, y, x_j * x_{j+1}, y, x_{j+2}, \dots, x_n) +$$

$$(16) \quad \sum_{i=j+2}^n \sum_{y \in R_q} (-1)^{i+3} (y, \dots, y, x_j * x_i, y, x_{j+1} * x_i, \dots, \widehat{x_i}, \dots, x_n)$$

In the above sums, $\widehat{x_i}$ denotes the omission of the element x_i . Notice, that we have the following cancellations:

$$\begin{aligned} (1) + (6) &= 0, \\ (5) + (13) &= 0, \\ (4) + (12) &= 0, \\ (10) + (16) &= 0, \\ (9) + (15) &= 0, \\ (11) + (14) &= 0. \end{aligned}$$

Moreover,

$$(2) + (3) = (-1)^j (f_n^j(x) - f_n^{j-1}(x)).$$

To finish the proof of the chain homotopy between $f_n^j(x)$ and $f_n^{j-1}(x)$, we need to look closer at

$$(7) = (-1)^{j+1} \sum_{y \in R_q} (y * x_j, \dots, y * x_j, y, x_{j+1}, \dots, x_n),$$

and

$$(8) = (-1)^{j+2} \sum_{y \in R_q} (y, \dots, y, x_j * y, x_{j+1}, \dots, x_n).$$

Notice, that

$$\sum_{y \in R_q} (y * x_j, \dots, y * x_j, y, x_{j+1}, \dots, x_n) = \sum_{y \in R_q} (y, \dots, y, y * x_j, x_{j+1}, \dots, x_n).$$

Therefore, in the case of R_3 , the commutativity property gives the final cancellation:

$$(7) + (8) = 0.$$

In the case of prime $q > 3$, we could consider maps $g_1^j, g_2^j: C_n^R(R_q) \rightarrow C_n^R(R_q)$ given by:

$$g_1^j(x) = \sum_{y \in R_q} (y, \dots, y, y * x_j, x_{j+1}, \dots, x_n),$$

and

$$g_2^j(x) = \sum_{y \in R_q} (y, \dots, y, x_j * y, x_{j+1}, \dots, x_n).$$

These are chain maps, if q is odd and not divisible by 3. The straightforward proof of this statement uses the fact that, for such q , the map $r_a: R_q \rightarrow R_q$ given by $r_a(b) = b * a * b$ is a bijection.

We conjecture the following:

Conjecture 4.3. *The maps g_1^j and $g_2^j: C_n^R(R_q) \rightarrow C_n^R(R_q)$ are chain homotopic, for prime $q > 3$.*

If this conjecture is true, then the maps $f_n^j(x)$ and $f_n^{j-1}(x)$ are chain homotopic for any odd prime q (and $j \geq 2$). This would allow us to prove the Conjecture 4.1. For now, we can finish the proof for $q = 3$. Recall that $\mathcal{O}(Q)$ denotes the set of orbits of a quandle Q . We can consider it as a trivial quandle ($a * b = a$, for any $a, b \in \mathcal{O}(Q)$). Let us define two maps, $\pi: C_n(R_q) \rightarrow C_n(\mathcal{O}(R_q))$, and $\Psi: C_n(\mathcal{O}(R_q)) \rightarrow C_n(R_q)$ in the following way:

$$\pi(x_1, \dots, x_n) = (1, \dots, 1),$$

where 1 denotes the single orbit of the quandle R_q ;

$$\Psi(1, \dots, 1) = \sum_{y \in R_q} (y, \dots, y).$$

Notice that both Ψ and π are chain maps. We have

$$\Psi\pi(x) = \sum_{y \in R_q} (y, \dots, y) = f_n^n(x).$$

Therefore, on homology, $\Psi_*\pi_* = q \cdot Id$. It follows that the torsion of $H_n^R(R_q)$ is annihilated by q , for $q = 3$. $\square$

5. $H_4^Q(R_p)$ IS NONTRIVIAL FOR p ODD PRIME

Theorem 5.1. $H_4^Q(R_p)$ contains Z_p , for p odd prime.

Proof. As we already know from Section 3,

$$\sum_{i=0}^{p-1} (0, i, i+1)$$

is a generator of $H_3^Q(R_p)$. Furthermore,

$$\alpha\left(\sum_{i=0}^{p-1} (0, i, i+1)\right)$$

is a generator of $H_3^R(R_p)$, because $\alpha_*: H_*^Q(R_p) \rightarrow H_*^R(R_p)$ is a monomorphism.

$$\Psi = \left(\alpha\left(\sum_{i=0}^{p-1} (0, i, i+1)\right), 0\right) + \left(\alpha\left(\sum_{i=0}^{p-1} (0, i, i+1)\right), 0\right) * 0$$

is nontrivial in $H_4^R(R_p)$ (the map $h'_0 = h_0 + (h_0) * a$ induces monomorphism on homology).

We will show that

$$\sum_{i=0}^{p-1} (0, i, i+1, 0) + \left(\sum_{i=0}^{p-1} (0, i, i+1, 0)\right) * 0$$

is nontrivial in $H_4^Q(R_p)$ by proving that

$$\begin{aligned}\Phi &= \alpha\left(\sum_{i=0}^{p-1}(0, i, i+1, 0) + \left(\sum_{i=0}^{p-1}(0, i, i+1, 0)\right) * 0\right) = \\ &\alpha\left(\sum_{i=0}^{p-1}(0, i, i+1, 0)\right) + \left(\alpha\left(\sum_{i=0}^{p-1}(0, i, i+1, 0)\right)\right) * 0\end{aligned}$$

is a cycle homologous to Ψ . That is, we prove that $\Phi - \Psi$ is zero in homology. Notice that

$$\alpha(0, i, i+1) = (0, i, i+1) - (0, 0, i+1) - (0, i, i) + (0, 0, i),$$

and therefore,

$$\alpha\left(\sum_{i=0}^{p-1}(0, i, i+1)\right) = \sum_{i=0}^{p-1}(0, i, i+1) - \sum_{i=0}^{p-1}(0, i, i).$$

Also,

$$\alpha(0, i, i+1, 0) = (\alpha(0, i, i+1), 0) - (\alpha(0, i, i+1), i+1);$$

thus,

$$\begin{aligned}\alpha\left(\sum_{i=0}^{p-1}(0, i, i+1, 0)\right) &= \sum_{i=0}^{p-1}(0, i, i+1, 0) - \sum_{i=0}^{p-1}(0, i, i, 0) \\ &\quad - \sum_{i=0}^{p-1}(0, i, i+1, i+1) + \sum_{i=0}^{p-1}(0, i, i, i+1).\end{aligned}$$

It follows that

$$\Phi - \Psi = \sum_{i=0}^{p-1}(0, i, i, i+1) - \sum_{i=0}^{p-1}(0, i, i+1, i+1) + \left(\sum_{i=0}^{p-1}(0, i, i, i+1) - \sum_{i=0}^{p-1}(0, i, i+1, i+1)\right) * 0.$$

Notice that

$$c = \sum_{i=0}^{p-1}(0, i, i, i+1) - \sum_{i=0}^{p-1}(0, i, i+1, i+1)$$

is a cycle homologous to

$$\left(\sum_{i=0}^{p-1}(0, i, i, i+1) - \sum_{i=0}^{p-1}(0, i, i+1, i+1)\right) * 0.$$

Therefore, to show that $(\Phi - \Psi) \sim 0$, it is enough to show that $c \sim 0$ (here, $\sim$ denotes homologous cycles).

It follows from the Proposition 2.1(iv) that

$$h'_{i+1}(h'_i(h'_i(0))) - h'_{i+1}(h'_{i+1}(h'_i(0))) \sim 0.$$

We have

$$\begin{aligned}h'_{i+1}(h'_i(h'_i(0))) &= h'_{i+1}(h'_i((0, i) + (0 * i, i))) = h'_{i+1}(2((0, i) + (0 * i, i))) = \\ &2((0, i, i, i+1) + (0 * i, i, i, i+1) + (0 * (i+1), i+2, i+2, i+1) + (0 * i * (i+1), i+2, i+2, i+1)),\end{aligned}$$

and

$$\begin{aligned} h'_{i+1}(h'_{i+1}(h'_i(0))) &= h'_{i+1}(h'_{i+1}((0, i) + (0 * i, i))) = \\ h'_{i+1}((0, i, i+1) + (0 * i, i, i+1) + (0 * (i+1), i+2, i+1) + (0 * i * (i+1), i+2, i+1)) &= \\ 2((0, i, i+1, i+1) + (0 * i, i, i+1, i+1) + (0 * (i+1), i+2, i+1, i+1) + (0 * i * (i+1), i+2, i+1, i+1)). \end{aligned}$$

From above, and the fact that there can be only odd torsion in homology of R_k with k odd, follows

$$\begin{aligned} & \left(\sum_{i=0}^{p-1} (0, i, i, i+1) + \sum_{i=0}^{p-1} (2i, i, i, i+1) + \sum_{i=0}^{p-1} (2i, i+1, i+1, i) + \sum_{i=0}^{p-1} (2, i+1, i+1, i) \right. \\ & \left. - \sum_{i=0}^{p-1} (0, i, i+1, i+1) - \sum_{i=0}^{p-1} (2i, i, i+1, i+1) - \sum_{i=0}^{p-1} (2i, i+1, i, i) - \sum_{i=0}^{p-1} (2, i+1, i, i) \right) \sim 0. \end{aligned}$$

The above expression decomposes into the following cycles:

$$(17) \quad c = \sum_{i=0}^{p-1} (0, i, i, i+1) - \sum_{i=0}^{p-1} (0, i, i+1, i+1),$$

$$(18) \quad \sum_{i=0}^{p-1} (2, i+1, i+1, i) - \sum_{i=0}^{p-1} (2, i+1, i, i),$$

$$(19) \quad \sum_{i=0}^{p-1} (2i, i, i, i+1) - \sum_{i=0}^{p-1} (2i, i+1, i, i),$$

$$(20) \quad \sum_{i=0}^{p-1} (2i, i+1, i+1, i) - \sum_{i=0}^{p-1} (2i, i, i+1, i+1).$$

We will show that all these cycles are homologous to c .

Let x be such that $2 * x = 0$. Then

$$\begin{aligned} (18) * x &= \left(\sum_{i=0}^{p-1} (2, i+1, i+1, i) - \sum_{i=0}^{p-1} (2, i+1, i, i) \right) * x = \\ \sum_{i=0}^{p-1} (0, 2x - (i+1), 2x - (i+1), 2x - i) &- \sum_{i=0}^{p-1} (0, 2x - (i+1), 2x - i, 2x - i) = \\ \sum_{i=0}^{p-1} (0, i, i, i+1) - \sum_{i=0}^{p-1} (0, i, i+1, i+1) &= (17). \end{aligned}$$

Thus, (17) $\sim$ (18).

Now, (17) - (19) is the following boundary:

$$\partial \left(\sum_{i=0}^{p-1} (0, i, i, i+1) + \sum_{i=0}^{p-1} (0, i, i+1, i+1) \right) =$$

$$\begin{aligned}
& \sum_{i=0}^{p-1} (0, i, i, i+1) - \sum_{i=0}^{p-1} (2i, i, i, i+1) - \sum_{i=0}^{p-1} (0, i, i, i) + \sum_{i=0}^{p-1} (2i+2, i+2, i+2, i+2) \\
& + \sum_{i=0}^{p-1} (0, i+1, i+1, i+1) - \sum_{i=0}^{p-1} (2i, i+1, i+1, i+1) \\
& - \sum_{i=0}^{p-1} (0, i, i+1, i+1) + \sum_{i=0}^{p-1} (2i+2, i+2, i+1, i+1) = \\
& \sum_{i=0}^{p-1} (0, i, i, i+1) - \sum_{i=0}^{p-1} (2i, i, i, i+1) - \sum_{i=0}^{p-1} (0, i, i+1, i+1) + \sum_{i=0}^{p-1} (2i, i+1, i, i) = (17) - (19).
\end{aligned}$$

Finally, we prove that (18) – (20) is a boundary.

$$\begin{aligned}
& \partial \left(\sum_{i=0}^{p-1} (2, i+1, i+1, i+1, i) + \sum_{i=0}^{p-1} (2, i+1, i, i, i) \right) = \\
& \sum_{i=0}^{p-1} (2, i+1, i+1, i) - \sum_{i=0}^{p-1} (2i, i+1, i+1, i) - \sum_{i=0}^{p-1} (2, i+1, i+1, i+1) + \sum_{i=0}^{p-1} (2i-2, i-1, i-1, i-1) \\
& + \sum_{i=0}^{p-1} (2, i, i, i) - \sum_{i=0}^{p-1} (2i, i, i, i) - \sum_{i=0}^{p-1} (2, i+1, i, i) + \sum_{i=0}^{p-1} (2i-2, i-1, i, i) = \\
& \sum_{i=0}^{p-1} (2, i+1, i+1, i) - \sum_{i=0}^{p-1} (2i, i+1, i+1, i) - \sum_{i=0}^{p-1} (2, i+1, i, i) + \sum_{i=0}^{p-1} (2i, i, i+1, i+1) = (18) - (20).
\end{aligned}$$

It follows that

$$4c \sim 0,$$

and this forces $c \sim 0$ (there can be no even torsion). Thus, $(\Phi - \Psi) \sim 2c \sim 0$, what ends the proof. $\square$

6. FUTURE DIRECTIONS

We have constructed, in Section 2, two homological operations $h'_0 : C_n^R(R_k) \rightarrow C_{n+1}^R(R_k)$ and $h_s : C_n^R(R_k) \rightarrow C_{n+2}^R(R_k)$. We have proven that h'_0 induces a monomorphism on rack homology $H_n^R(R_k) \rightarrow H_{n+1}^R(R_k)$ for k odd. We have conjectured that $(h_s)_* : H_n^R(R_k) \rightarrow H_{n+2}^R(R_k)$ is also a monomorphism. More generally, we propose:

Conjecture 6.1. $(h'_0)_* \oplus (h_s)_* : H_{n+1}^R(R_k) \oplus H_n^R(R_k) \rightarrow H_{n+2}^R(R_k)$ is a monomorphism for k odd and $n > 1$.

In the case of quandle homology, the composition $h'_a h'_b$ induces the zero map on homology for any $a, b \in R_k$. Therefore, instead of this map, we consider the composition map $h'_{(s,0)} = h'_0 h_s : C_n^Q(R_k) \rightarrow C_{n+3}^Q(R_k)$ and propose:

Conjecture 6.2. $(h_s)_* \oplus (h'_{(s,0)})_* : H_{n+1}^Q(R_k) \oplus H_n^Q(R_k) \rightarrow H_{n+3}^Q(R_k)$ is a monomorphism for k odd and $n > 1$.

We have checked that the conjecture holds for $k = 3, 5$ and $n = 2, 3$ (in these cases, these maps are isomorphisms) and for $k = 3$ and $n = 4, 5$.

Conjecture 6.2 is not sufficient to prove our main conjecture (Conjecture 1.5). To achieve this, we need additional homological operation of degree 4: $h_{new} : C_n^Q(R_k) \rightarrow C_{n+4}^Q(R_k)$ such that

$$(h_s)_* \oplus (h'_{(s,0)})_* \oplus (h_{new})_* : H_{n+2}^Q(R_k) \oplus H_{n+1}^Q(R_k) \oplus H_n^Q(R_k) \rightarrow H_{n+4}^Q(R_k)$$

is an isomorphism for k odd and $n > 1$. It would provide inductive step in a proof of Conjecture 1.5, as $f_{n+4} = f_{n+3} + f_{n+1} = f_{n+2} + f_{n+1} + f_n$.

7. APPENDIX 1

In this section we demonstrate a generalization of Corollary 2.4.

Theorem 7.1. *Let a quandle Q satisfy the quasigroup property (Definition 2.2). Then, if N annihilates the torsion of $H_n^R(Q)$, then $|Q|N$ annihilates the torsion of $H_{n+1}^R(Q)$.*

Proof. Consider the composition of two homomorphisms

$$gf : C_n^R(Q) \rightarrow C_{n-1}^R(Q) \rightarrow C_n^R(Q),$$

where

$$f(x_1, x_2, \dots, x_n) = (-1)^n(x_2, \dots, x_n)$$

and

$$g(x_2, \dots, x_n) = (-1)^n \sum_{y \in Q} (y, x_2, \dots, x_n).$$

We check that f and g are chain maps.

$$\begin{aligned} fd(x_1, x_2, \dots, x_n) &= \\ (-1)^{n-1} \sum_{i=2}^n (-1)^i ((x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n) - (x_2 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)) &= \\ df(x_1, \dots, x_n). \\ dg(x_2, \dots, x_n) &= \\ (-1)^n \sum_{y \in Q} \sum_{i=2}^n (-1)^i ((y, \dots, x_{i-1}, x_{i+1}, \dots, x_n) - (y * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)) &= \\ (-1)^n \sum_{y \in Q} \sum_{i=3}^n (-1)^i ((y, \dots, x_{i-1}, x_{i+1}, \dots, x_n) - (y, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)) &= \\ g(\sum_{i=3}^n (-1)^{i+1} ((x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n) - (x_2 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n))) &= \\ gd(x_2, \dots, x_n). \end{aligned}$$

Furthermore, as shown in [L-N](Section 3), the composition gf is chain homotopic to $|Q|Id$. Therefore, on homology, $(gf)_* = |Q|Id$. On the other hand, $(gf)_* = g_*f_*$. Thus, if $Nf_*(a) = 0$, then $0 = Ng_*f_*(a) = N|Q|a$, and the theorem follows. $\square$

8. APPENDIX 2

In this section, we work exclusively with quandle homology, and we show that for R_k , k odd, we can compute homology from a smaller chain complex of “symmetric” elements.

We will work with homology with coefficients in any ring $\mathcal{R}$ (treated as an abelian group). Let $C_n^{inv(a)}(Q; \mathcal{R})$ denote the subgroup of $C_n(Q; \mathcal{R})$ composed of elements w which satisfy $w = w * a$. $C_*^{inv(a)}(Q; \mathcal{R})$ is a subchain complex of $C_*(Q; \mathcal{R})$. Indeed, $*_a(dw) = d(*_a w) = dw$.

Let $C_n^{cov(a)}(Q; \mathcal{R}) = C_n(Q; \mathcal{R}) / C_n^{inv(a)}(Q; \mathcal{R})$. We have a short exact sequence of chain complexes:

$$0 \rightarrow C_*^{inv(a)}(Q; \mathcal{R}) \xrightarrow{i} C_*(Q; \mathcal{R}) \rightarrow C_*^{cov(a)}(Q; \mathcal{R}) \rightarrow 0,$$

and related long exact sequence of homology

$$\begin{aligned} \dots \rightarrow H_{n+1}^{cov(a)}(Q; \mathcal{R}) &\xrightarrow{\partial} H_n^{inv(a)}(Q; \mathcal{R}) \xrightarrow{i_*} H_n(Q; \mathcal{R}) \rightarrow \\ &\rightarrow H_n^{cov(a)}(Q; \mathcal{R}) \xrightarrow{\partial} H_{n-1}^{inv(a)}(Q; \mathcal{R}) \xrightarrow{i_*} H_{n-1}(Q; \mathcal{R}) \rightarrow \dots \end{aligned}$$

Proposition 8.1. *The map h_a restricted to $C_n^{inv(a)}(Q; \mathcal{R})$ is a chain map.*

Proof. $dh_a(w) = h_a d(w) + (-1)^n (w - w * a) = dh_a(w)$, as $w \in C_n^{inv(a)}(Q; \mathcal{R})$. $\square$

Lemma 8.2. *If $\mathcal{R}$ is a ring with 2 invertible, and Q is a kei, then the above short exact sequence of chain complexes splits. In particular, the connecting homomorphism ∂ , in the long exact sequence of homology, is a zero map.*

Proof. Let $\{C_n^+\}$ be a subchain complex of $\{C_n(Q; \mathcal{R})\}$ generated by the symmetric elements of the form $u + u * a$, where $u \in Q^n$. Similarly, let $\{C_n^-\}$ be a subchain complex of $\{C_n(Q; \mathcal{R})\}$ generated by the antisymmetric elements $u - u * a$. Notice that $*_a(w) = w$ for $w \in C_n^+$, and $*_a(v) = -v$ for $v \in C_n^-$. It follows that $C_n^+ \cap C_n^- = 0$. Indeed, if $w \in C_n^+ \cap C_n^-$, then $*_a(w) = w$ and $*_a(w) = -w$, therefore, $w = 0$ (C_n^+ and C_n^- are $\mathcal{R}$ -modules with 2 invertible in $\mathcal{R}$).

Furthermore, $C_n(Q; \mathcal{R}) = C_n^+ \oplus C_n^-$, because for any element $w \in C_n(Q; \mathcal{R})$, we have $w + w * a + w - w * a \in C_n^+ \oplus C_n^-$, so $2w \in C_n^+ \oplus C_n^-$. Since 2 is invertible in $\mathcal{R}$, $w \in C_n^+ \oplus C_n^-$.

Thus, the above short exact sequence of chain complexes splits. From the fact that $C_n^+ \subset C_n^{inv(a)}(Q; \mathcal{R})$ and $C_n^- \cap C_n^{inv(a)}(Q; \mathcal{R}) = 0$, follows that $C_n^+ = C_n^{inv(a)}(Q; \mathcal{R})$. Similarly, $C_n^- = C_n^{cov(a)}(Q; \mathcal{R})$. $\square$

Lemma 8.3. *If Q is a kei, and the homomorphism $2Id : H_n(Q; \mathcal{R}) \rightarrow H_n(Q; \mathcal{R})$ is an isomorphism (e.g., H_n is a finite abelian group without even torsion, or $\mathbb{Z}[\frac{1}{2}]$ -module), then the homomorphism $i_* : H_n^{inv(a)}(Q; \mathcal{R}) \rightarrow H_n(Q; \mathcal{R})$ is an epimorphism.*

Proof. If the cycles $w_1, \dots, w_s$ are generators of $H_n(Q; \mathcal{R})$, then also $2w_1, \dots, 2w_s$ generate this group. By Proposition 2.1(i), $w + w * a = 2w$ in $H_n(Q; \mathcal{R})$, therefore, a -symmetric elements generate $H_n(Q; \mathcal{R})$, and the lemma follows. $\square$

As a consequence, we have the result which says that for odd dihedral quandles R_k , the homology $H_n^{inv(a)}(R_k)$ determines the quandle homology $H_n(R_k)$. More precisely:

Theorem 8.4. *For $n > 1$, the homomorphism $H_n^{inv(a)}(R_k) \rightarrow H_n(R_k)$, yielded by embedding of chain complexes, is an epimorphism, with the kernel being a 2-torsion part of $H_n^{inv(a)}(R_k)$.*

Proof. We use Lemmas 8.2 and 8.3 for homology of R_k , and the ring $\mathcal{R} = \mathbb{Z}[\frac{1}{2}]$. We get the isomorphism:

$$H_n^{inv(a)}(R_k; \mathbb{Z}[\frac{1}{2}]) \rightarrow H_n(R_k; \mathbb{Z}[\frac{1}{2}]).$$

From the universal coefficient theorem for homology, and the fact that $\mathbb{Z}[\frac{1}{2}]$ is a torsion free group, we get the isomorphism

$$H_n^{inv(a)}(R_k) \otimes \mathbb{Z}[\frac{1}{2}] \rightarrow H_n(R_k) \otimes \mathbb{Z}[\frac{1}{2}].$$

The theorem follows, because $H_n(R_k)$ is a group without a 2-torsion [L-N]. $\square$

REFERENCES

- [CJKLS] S. Carter, D. Jelsovsky, S. Kamada, L. Langford, M. Saito, State-sum invariants of knotted curves and surfaces from quandle cohomology, *Electron. Res. Announc. Amer. Math. Soc.* **5**, 1999, 146-156;
- [CJKS] S. Carter, D. Jelsovsky, S. Kamada, M. Saito, Quandle homology groups, their Betti numbers, and virtual knots, *J. Pure Appl. Algebra* **157**, 2001, 135-155;
e-print: <http://front.math.ucdavis.edu/math.GT/9909161>
- [CKS] S. Carter, S. Kamada, M. Saito, Surfaces in 4-space, *Encyclopaedia of Mathematical Sciences, Low-Dimensional Topology III*, R.V.Gamkrelidze, V.A.Vassiliev, Eds., 213pp.
- [E-G] P. Etingof, M. Grana, On rack cohomology, *J. Pure Appl. Algebra* **177**, 2003, 49-59;
e-print: <http://front.math.ucdavis.edu/math.QA/0201290>
- [F-R] R. Fenn, C. Rourke, Racks and links in codimension two, *Journal of Knot Theory and its Ramifications* **1**(4), 1992, 343-406;
e-print: <http://www.maths.sussex.ac.uk/Staff/RAF/Maths/racks.ps>
- [FRS] R. Fenn, C. Rourke and B. Sanderson, James bundles and applications, preprint;
e-print: <http://www.maths.warwick.ac.uk/~cpr/ftp/james.ps>
- [GAP4] The GAP Group, GAP – Groups, Algorithms, and Programming,
<http://www.gap-system.org>;
- [Ha] A. Hatcher, Algebraic Topology, Cambridge University Press, 2002;
<http://www.math.cornell.edu/~hatcher/AT/ATch3.pdf>
- [Joy] D. Joyce, A classifying invariant of knots: the knot quandle, *J. Pure Appl. Algebra* **23**, 1982, 37-65;

- [L-N] R. A. Litherland, S. Nelson, The Betti numbers of some finite racks, *J. Pure Appl. Algebra* **178**, 2003, 187-202;
e-print: <http://front.math.ucdavis.edu/math.GT/0106165>
- [Moc] T. Mochizuki, Some calculations of cohomology groups of finite Alexander quandles, *J. Pure Appl. Algebra* **179**, 2003, 287-330;
- [N-P] M. Niebrzydowski, J. H. Przytycki, Burnside Kei, *Fundamenta Mathematicae* **190**, June, 2006, 211-229;
e-print: <http://front.math.ucdavis.edu/math.GT/0601004>
- [Oht] T. Ohtsuki, Quandles, in Problems on invariants of knots and 3-manifolds, *Geometry and Topology Monographs*, Volume 4, 2003, 455-465;
e-print: <http://front.math.ucdavis.edu/math.GT/0406190>
- [Tak] M. Takasaki, Abstraction of symmetric transformation, (in Japanese) *Tohoku Math. J.* **49**, 1942/3, 145-207; the English translation is being prepared by S. Kamada, to be published in Proceedings of Knots in Washington XX; 60th birthday of Louis H. Kauffman.

Maciej Niebrzydowski
e-mail: mniebrz@gmail.com

Józef H. Przytycki
e-mail: przytyck@gwu.edu

Józef Przytycki is on sabbatical leave at University of Maryland, College Park.

DEPT. OF MATHEMATICS, OLD MAIN BLDG., 1922 F ST. NW
THE GEORGE WASHINGTON UNIVERSITY, WASHINGTON DC, 20052